

EFI-98-24

UT-KOMABA/98-15

June 1998

# Short-Distance Space-Time Structure and Black Holes in String Theory : A Short Review of the Present Status \*

MIAO LI <sup>†</sup>

*Enrico Fermi Institute, University of Chicago  
5640 Ellis Avenue, Chicago, IL 60637*

TAMIAKI YONEYA <sup>‡</sup>

*Institute of Physics, University of Tokyo  
Komaba, Meguro-ku, 153 Tokyo*

## Abstract

We briefly review the present status of string theory from the viewpoint of its implications on the short-distance space-time structure and black hole physics. Special emphases are given on two closely related issues in recent developments towards nonperturbative string theory, namely, the role of the space-time uncertainty relation as a qualitative but universal characterization of the short-distance structure of string theory and the microscopic formulation of black-hole entropies. We will also suggest that the space-time uncertainty relation can be an underlying principle for the holographic property of M theory, by showing that the space-time uncertainty relation naturally explains the UV/IR relation used in a recent derivation of the holographic bound for D3 brane by Susskind and Witten.

---

\* To appear in the special issue of the *Journal of Chaos, Solitons and Fractals* on "Superstrings, M,F,S...Theory".

<sup>†</sup> e-mail address: [mli@curie.uchicago.edu](mailto:mli@curie.uchicago.edu)

<sup>‡</sup> e-mail address: [tam@hepl.c.u-tokyo.ac.jp](mailto:tam@hepl.c.u-tokyo.ac.jp)

## 1. Historical introduction

String theory emerged originally from the discovery of the Veneziano formula [1] for the scattering amplitude for mesons, satisfying the so-called  $s$ - $t$  duality in an idealized form. The  $s$ - $t$  duality says that the scattering amplitude can be expressed in two equivalent ways, either as the sum of resonances in the  $s$  channel or the particle exchanges in the  $t$  channel. However, it soon turned out, through various developments in extending the formula, that the requirement of the exact  $s$ - $t$  duality was too restrictive for the theory of strong interactions. The latter is then established as the gauge field theory of ‘color’ degrees of freedom, *Quantum Chromo Dynamics*.

On the other hand, it also turned out that there existed astonishingly beautiful and rich structure behind the formula [2]. In particular, the basic dynamical degrees of freedom governing the amplitude was identified [3] to be one-dimensionally extended objects, *strings*, instead of particles in ordinary local field theories. One of the most important aspects of the string picture was to make it possible to identify the symmetry ensuring the exact  $s$ - $t$  duality as the conformal symmetry in the dynamics of strings. Namely, the world-sheets formed as the trajectory of strings in space-time behave as Riemann surfaces. Most of the restrictive features of the string theory come from the requirement of the (super) conformal symmetry.

Here are some examples from the list of the properties of strings which result from the conformal symmetry: The critical dimensions, namely the conditions for the dimensionality of space-times such as 26 in bosonic models and 10 in the models with fermionic strings, are the conditions for cancellation of the conformal anomaly which usually plagues any quantum field theories to preserve the conformal symmetry of classical theory. The fixed-point equation in the renormalization-group formulation of the world-sheet dynamics contains the Einstein equations for background space-time in the long-distance regime. Cancellations of the ultra-violet divergences associated with the loop diagrams of string world sheets, one-to-one correspondence between the spectrum and the allowed external fields for the propagation of strings, .. and so on, are also the direct consequences of the conformal symmetry.

All of these properties are crucial ingredients which constitute the bases for the in-

terpretation of string theory as the unified quantum theory of all interactions including gravity. In particular, the string theory showed us, for the first time, that there existed a natural avenue towards the quantum theory of gravity by circumventing [4] the non-renormalizability of the ultra-violet divergences, since it turned out that the spectrum of consistent string theories necessarily contain the graviton whose interaction in the long-distance regime is correctly described [5][6] by the classical general relativity, adjoined with some additional fields like dilaton, antisymmetric tensor fields and further fermionic fields (called gravitinos) associated with space-time supersymmetry [7], namely, supergravity.

However, the present formulations of string theories are far from satisfactory. The string picture only provides us a set of perturbative rules for constructing the S-matrices in the expansion with respect to the strength of the string coupling  $g_s$  which is related to the vacuum expectation value of the dilaton field  $\phi$  as  $g_s = e^\phi$ . That is, we have seen only the surface of string theory in the special asymptotic limit  $\phi \rightarrow -\infty$ . From experiences in gauge-field theories for unifying the other interactions than gravity and the role of the non-perturbative behaviors in QCD, we cannot hope that such a perturbation expansion alone is sufficient to explore the physics near Planck scale where the very dynamics of the vacua and their transitions must be investigated. Present perturbative formulation has no power in giving definite predictions in such regime. All what we could envisage from the perturbative string theory is that there must exist certain tremendous theoretical structure whose real meaning and significance are not yet fully understood.

The developments [8] occurred in recent few years are very encouraging in this respect. Firstly, various duality relations which connect different perturbative string theories were discovered, and passed many nontrivial checks. The T-duality relates different background fields by the duality transformation on the world sheet of strings. For example, a string theory with a space-time dimension which is compactified into a circle with a radius  $R$  is equivalent with a string theory with the inverse compactified radius  $\ell_s/R$ , where  $\ell_s$  is the length constant characterizing the space-time extension of strings [9]. The S-duality [10], on the other hand, interchanging the roles of the conserved Ramond-Ramond and Neveu-Schwarz charges, relates a string theory with a given string coupling  $g_s$  to a string theory with inverse string coupling  $1/g_s$ . Furthermore, it turned out that, in addition to the pre-

viously known perturbative string theories, there must exist a new theory, called M-theory, whose perturbative behavior in the long-distance limit is described by 11 dimensional supergravity. From the viewpoint of M-theory [11], the type IIA string perturbation theory is a reduced model in which one of the spatial dimension is compactified with the radius  $R = g_s \ell_s$ . Hence, in the perturbative regime  $g_s \sim 0$ , the effective space-time dimension is 10.

In all these developments, one of the most crucial aspects is the discovery of the role of Dirichlet branes [12] (usually called ‘D-branes’) which carry the Ramond-Ramond charges. The elementary string states can never carry the Ramond-Ramond charges. Thus D-branes made it possible to discuss the properties of S-dual transformed theories in concrete forms.

Furthermore, D-branes opened up entirely new viewpoints for string theory. It has been known that there are a large number of classical solutions [13] for the low-energy effective field theories corresponding to the perturbative string theories, which contain various types of black-hole configurations. From the viewpoint of D-branes, most of them can be regarded as composite systems of D-branes. Once such identifications are made, we can count the number of physical states with given mass and charges, at least in the limit of extremal black holes where the masses of black holes approach to the lower bounds proportional to the charges. The latter limit corresponds to limits (called the BPS limit) where the space-time supersymmetry is partially recovered for the D-brane configurations. The results of counting, obtained by neglecting the string interactions, are reliable owing to the protection of supersymmetry. In this way, Bekenstein-Hawking’s classical formula for the black-hole entropy was derived [14] [15], for the first time, by the quantum-statistical method. In view of an analogy with the role played by black-body radiation in the early development of quantum theory in the beginning of this century, the statistical derivation of the black hole entropy should be regarded as an important testing ground where we can check whether or not any theory of quantum gravity correctly identifies the basic microscopic degrees of freedom. Thus string theory now turned out to be really the qualified candidate for the theory of quantum gravity from the viewpoint of black hole physics too. Not only that, these achievements clearly indicate that the space-time structure near Planck length or shorter should be interpreted in a drastically

different way from the classical geometrical picture, as has been emphasized in many occasions from various different perspectives.

Another important new aspect of string theory suggested from the advent of D-branes is the possibility of reconstructing string theory by using the D-branes as fundamental degrees of freedom. In other words, the original string may now be interpreted as a sort of collective modes in such a formulation. The hope in doing this is that such a reformulation may well be suitable to explore the non-perturbative regime of string theory and, more ambitiously, to obtain the rigorous non-perturbative definition of the theory.

In the usual perturbative string theory, the D-branes are treated by introducing open strings with Dirichlet boundary conditions (hence the name ‘Dirichlet’ brane), which impose the positions of the D-branes to be the ends of the open strings. Then the positions of the D-branes can be regarded as collective coordinates of string theory. The low-energy effective theory for such collective coordinates [16] are the super Yang-Mills theory, whose interpretation is of course entirely different from the usual one as for the unified theory of forces other than the gravity. Namely, the theory lives on the world volume of the D-branes, and the diagonal parts of the Higgs fields now play the role of transverse coordinates of the D-branes. All of other degrees of freedom, i.e. off-diagonal components and the gauge fields, are supposed to describe the dynamics of short open strings interacting with D-branes through the Dirichlet condition. Attempts towards the reconstruction of string theory are based on these effective super Yang-Mills theories. The first attempt along this line, so-called Matrix theory [17], is to hypothesize the lowest dimensional D-brane in the type IIA theory, zero-dimensional branes=D-particles, as the fundamental building blocks, or partons, of the theory and the effective super Yang-Mills theory with 9 Higgs fields, i.e. 9 dimensions, to be the exact theory in the infinite momentum frame (IFM) in 11 dimensional space-time. Another attempt called the type IIB matrix model [18], followed after this proposal, is to identify the zero-dimensional super Yang-Mills theory with 10 Higgs fields, namely, just the dimensional reduction of the 10D super Yang-Mills to a point, as the underlying non-perturbative theory for the type IIB strings in 10 dimensional space-time. In both cases, taking the large  $N$  limit is assumed to be an essential requirement for the exact nonperturbative formulation of the theory. In the former case, there is some indication that the theory is meaningful even for

finite  $N$  in the viewpoint of the discrete light-cone quantization [19][20]. Although there is no definitive verification that these proposals can be accepted as the correct definitions of the nonperturbative string theory or M theory, we already have reasonable amount of evidence for believing that these are indeed promising starting points, at the least, towards our goal.

Besides various technical problems, one of the mysteries associated with these proposals is what the underlying principle which selects the super Yang-Mills theory is, and why such Yang-Mills theory, provided it is the correct starting point, could give rise to the consistent theory of all interactions including gravity. For example, we do not know definite symmetry structure in super Yang-Mills theory which ensures the emergence of General Relativity in the long-distance regime, except for the existence of maximal supersymmetry which is only global from the viewpoint of the target space-time. It is not at all clear whether the maximal supersymmetry is sufficient to guarantee the emergence of consistent (namely, finite and unitary) quantum theory of gravity, beyond its role in effective low-energy descriptions of near BPS dynamics of D-branes. For example, we do not know any systematic method for extending the matrix models to general backgrounds, since we do not have a firm understanding on what the degrees of freedom in the super Yang-Mills descriptions representing the backgrounds are. In the case of Matrix theory, in particular, how to formulate the theory in Lorentz invariant fashion remains as a big mystery. In the case of the type IIB matrix model, although Lorentz covariant by its definition, we do not know how to discuss the geometry of backgrounds fields even in the long-distance approximation, and hence we cannot discuss the symmetry under the S-duality transformation which is expected to be the exact symmetry of the theory. These difficulties clearly indicate that some crucial things are missing in our present understanding of the matrix model approaches towards non-perturbative string or M theory.

Thus it seems to us that one of the most important tasks now is to seek for the definite guiding principles which explain the emergence of Yang-Mills theory and on the bases of which we can proceed toward more satisfactory formulation and definition of the theory. In the following, we review some of the attempts along this line of thoughts from two specific viewpoints, namely, the space-time uncertainty relation as a possible universal characterization of the short-distance space-time structure in string theory, and

the possibility of microscopic theory of black holes from the viewpoint of D-brane matrix models. The reason why we emphasize these two viewpoints must now be clear. Compared with classical General Relativity and gauge-quantum field theories of other interactions than gravity, the unified theory should be characterized by its departure from the former with respect to the short-distance space-time structure and the fundamental microscopic degrees of freedom at the shortest possible space-time scale.

In section 2, we motivate and summarize the space-time uncertainty principle [21] and discuss its roles in the dynamics of D-branes, especially in the case of D-particles. In section 3, we briefly review the entropy problem of black holes and discuss the statistical derivation of entropy from Matrix-theory viewpoint. In section 4, we will suggest a generalized space-time uncertainty relation suitable for M theory and conclude by pointing out that the space-time uncertainty relation explains some of the crucial properties leading to the holographic bound for the entropy of D3 brane.

## 2. Space-time uncertainty principle of string theory

### *2.1 World-sheet conformal symmetry and the short-distance space-time structure*

We have emphasized that the symmetry which leads to the key features of string theory as the unified theory of all interactions is the conformal symmetry exhibited in the dynamics of world sheets of strings. It is therefore reasonable to expect that the conformal symmetry must be taken into account in the nonperturbative reformulation of the theory. However, the conformal symmetry as the symmetry of the world-sheet dynamics seems to be intrinsically a perturbative concept, depending upon the emergence of Riemann surfaces in the perturbative formulation. The Riemann surfaces in string theory, or more precisely, their equivalence classes called the moduli space of Riemann surfaces, replace the role of the space of proper times for relativistic point particles. The proper time is intrinsically a perturbative concept in quantum field theory. It is therefore important to reinterpret the conformal symmetry using physical quantities which do not intrinsically depend upon the perturbation theory.

The importance of reinterpreting the conformal symmetry is also expected from the following consideration. The world-sheet dynamics of perturbative string theory is con-

structed using the conformal field theories (CFTs). From the general viewpoint of two-dimensional quantum field theories, CFTs are a very special class of field theories, characterized by the fixed-points of renormalization-group equation. In other words, the astonishingly desirable features and consistency of string theory as the unified theory are realized only by choosing a special class of two-dimensional field theories describing the dynamics of world sheets. We can point out an amusing analogy here. The requirement of conformal symmetry is playing a somewhat analogous role with the Bohr-Sommerfeld quantization condition for the adiabatic invariants in classical analytical dynamics. The consistency of imposing the latter condition in classical theory comes from the adiabatic theorem which says that the adiabatic invariants remain invariant under slow changes of external variables. Hence the quantization condition on them, although ad hoc from the viewpoint of the continuous degrees of freedom of possible initial conditions, can be protected against small perturbations. The analog to the adiabatic theorem in quantum field theories is the fixed-point condition of renormalization group. Of course, by the advent of quantum mechanics, the ad hoc quantization conditions were replaced by the operator algebra of physical observables which are characterized by canonical commutation relations. Similarly, we should perhaps expect that the requirement of conformal symmetry in the perturbative formulation of string theory is replaced by some more general condition which can be expressed without using the intrinsic properties of rules for constructing string S-matrices.

With these motivations in mind, one of the present authors proposed to reformulate the world-sheet conformal symmetry as a simple uncertainty relation in space-time [22]. Let us first briefly recapitulate the space-time uncertainty relation.<sup>§</sup>

Consider a parallelogram on the string world-sheet and the Polyakov amplitude for the mapping from the world-sheet to a region of the target space-time. Let the lengths of the two orthogonal sides in the world-sheet parallelogram be  $a, b$  in the conformal gauge where  $\dot{x} \cdot x' = 0$ ,  $\dot{x}^2 + x'^2 = 0$  and the corresponding physical space-time length be  $A, B$ , respectively. Then apart from the power behaved pre-factor, the amplitude is proportional

---

<sup>§</sup> Independently of the proposal of the space-time uncertainty relation, an intimately related concept of the ‘minimal length’ was suggested by several authors [23][24] and has got popularity [25]. It should be noted that the space-time uncertainty relation is more general than the concept of the minimal length which now turned out to be restricted to the perturbative string amplitudes.



to

$$\exp\left[-\frac{1}{\ell_s^2}\left(\frac{A^2}{\Gamma} + \frac{B^2}{\Gamma^*}\right)\right] \quad (2.1)$$

where

$$\Gamma \equiv \frac{a}{b}, \quad \Gamma^* \equiv \frac{b}{a}, \quad (\Gamma\Gamma^* = 1). \quad (2.2)$$

Due to the conformal invariance, the amplitude depends on the Riemann sheet parameters only through the ratio  $\Gamma$  or  $\Gamma^*$ , which are called the *extremal length* and the *conjugate extremal length*, respectively. Clearly, the relation  $\Gamma\Gamma^* = 1$  leads to the uncertainty relation [22][26]

$$\Delta T \Delta X \sim \ell_s^2 \sim \alpha' \quad (2.3)$$

taking the  $A$  direction to be time-like  $\Delta T \sim \langle A \rangle$  and hence the  $B$  direction to be space-like  $\Delta X \sim \langle B \rangle$ . The obvious relation  $\Gamma\Gamma^* = 1$  is the origin of the familiar modular invariance of the torus amplitudes in string theory. The extremal length is the most fundamental moduli parameter characterizing conformal invariants, in general. Since arbitrary amplitudes can be constructed by pasting together many parallelograms, any string amplitudes satisfy the above reciprocal relation qualitatively. Although this form looks too simple as the characterization of the conformal invariance, it has a virtue that its validity is very general, as we will explain shortly, and does not use the concepts which depend intrinsically on perturbation theory. Our proposal [26][27] is to use this relation as one of possible guiding principles towards nonperturbative reformulation of string theory and M-theory.

We note that the above uncertainty relation suggests that the classical space-time is something very analogous to the classical phase space of particle mechanics. Indeed the form (2.1) of the amplitude is reminiscent of the Wigner representation of the Gaussian wave packet of a point particle. In this sense, our proposal is similar to other proposals [29] of space-time quantization. Most of such proposals, however, presuppose the existence of minimum length in all space-time directions. Our proposal is quite different in this respect, since the above relation does not prohibit the existence of shorter lengths than the string scale  $\ell_s$  in a particular direction. It only says that there is a duality between short and large distance phenomena in string theory. The importance of shorter length

scales from the more recent context of various duality relations in string theory has been emphasized by Shenker in ref. [32].

The uncertainty relation (2.3) is consistent with an elementary property of strings that the energy of a string is roughly proportional to its space-time length.

$$\Delta E \sim \frac{\hbar}{\alpha'} \Delta X_l. \quad (2.4)$$

with  $X_l$  being the length of a string measured along its longitudinal direction. Then the ordinary time-energy uncertainty relation  $\Delta T \Delta E \geq \hbar$  leads to (2.3) [22]. It is important here to discriminate the length scales in the longitudinal and transverse directions with respect to the string. As is well known, the (squared) transverse length scale grows logarithmically with energy used to probe the strings. This explains the linearly rising Regge trajectory for the Regge-pole behavior in high-energy peripheral scattering. ¶ The dual role of the time and the longitudinal spatial lengths is a natural space-time expression of the original  $s$ - $t$  duality. The particle exchange (Regge exchange in the old language) and the resonance behaviors correspond to the regimes,  $\Delta X_l \rightarrow \infty$  and  $\Delta T \rightarrow \infty$ , respectively. Furthermore, the Regge behavior is consistent with the existence of graviton, since scattering amplitudes in general are expected to be roughly proportional to  $\Delta X_l \sim \ell_s^2 / \Delta T \propto E$  which implies, by adopting the argument in [33], that the intercept  $\alpha(0)$  of the leading Regge trajectory is 2, from the relation  $E \sim E^{\alpha(t)-1}$ .

On the other hand, in the high energy fixed-angle scatterings with large  $s$ - and  $t$ -channel momenta studied in detail in [24], we are trying to probe the region where both the time and the spatial scales are small. Clearly, such a region is incompatible with the space-time uncertainty relation. The exponential fall-off of the perturbative string amplitudes in this limit may be interpreted as a manifestation of this property. According to the space-time uncertainty relation, at least one of the two length scales,  $\Delta X$  or  $\Delta T$ , must be larger than the string scale  $\ell_s$ . Therefore there is no degrees of freedom left in this regime. It is well known that any consistent theory of quantum gravity must indicate lessening of the

---

¶Note that in the special case of the strings in 1+1 dimensions, there is a longitudinal size but no transverse size, and, hence, the high-energy limits of the perturbative amplitudes show power-law behaviors, apart from the phase factors associated with the external legs. For a discussion on the physics of the 1+1 dimensional string scattering, we refer the reader to ref. [30]. For a review on the connection of 1+1 dimensional strings and old-fashioned matrix models, see [31] and references therein.

degrees of freedom near the Planck scale where the quantum nature of gravity becomes important. We have seen that the space-time uncertainty relation can indeed be a natural mechanism for this. Qualitatively, it is also consistent with the known high-temperature behavior [28] of the perturbative string amplitude, since the high-temperature limit is effectively equivalent to considering the limit  $\Delta T \rightarrow 0$ .

## 2.2 Black hole physics and the space-time uncertainty relation

The space-time uncertainty relation is also expected to be important in black hole physics, since black hole provides us a natural tool for probing the short time region. By definition, the black hole horizon is the region where, owing to infinite red shift effect, any finite time interval in the asymptotic space-time boundary corresponds to infinitely short time interval  $\Delta T \sim 0 \rightarrow \Delta X \rightarrow \infty$ . Thus we expect that the black hole scatterings can be a unique arena where we may check the stringy behavior of the theory. Perhaps the black hole horizon is one of the places where the predictions and interpretations of string theory could significantly deviate from those of local field theory, since there we cannot rely upon the usual intuition based on familiar local field theory. For example, some peculiar behaviors found for a matrix model for the 1+1 dimensional black hole [34] might be due to this. For a review on this controversial issue, we refer the reader to [35] and the references therein. Unfortunately, most of the reliable discussions of the black-hole scattering of strings are based on the low-energy limit ( $\alpha' \sim 0$ ). However, the recent developments of the D-brane technology and Matrix theory indeed suggest a picture of the black-hole scattering which is quite different from the local field theory with respect to the puzzle of information loss [36].

Qualitatively, the deviation from the local field theory and string theory should manifest itself when the length scale of black holes becomes comparable to the string scale where the space-time uncertainty relation becomes essential. There is indeed a signature that the stringy nature becomes dominant beyond this regime. In D-dimensional space-time, the Bekenstein-Hawking entropy of a Schwarzschild black hole of mass  $E$  is given by

$$S \left( = \frac{\text{Area}}{4G_N} \right) = \frac{1}{4} G_N^{\frac{1}{D-3}} E^{\frac{D-2}{D-3}}. \quad (2.5)$$

On the other hand, the massive (super) string state has a quantum degeneracy of the

order

$$d(E) = \exp 2\pi \sqrt{\frac{D-2}{6}} N \quad (2.6)$$

with  $N = (\ell_s E)^2$ . If we identify the Schwarzschild black hole of string size  $\ell_s$  with a massive string state by setting <sup>||</sup>

$$\ell_s \rightsquigarrow \ell_{\text{eff}} \sim R_{\text{Schwarzschild}}, \sim (G_N E)^{\frac{1}{D-3}},$$

we have

$$\log d(N) \sim G_N^{\frac{1}{D-3}} E^{\frac{D-2}{D-3}} \sim S_{\text{BH}}. \quad (2.7)$$

This strongly suggests that the black hole with sizes comparable with string scale must be treated by string theory, and the latter would possibly provide the microscopic explanation of the entropy. More detailed examination of the correspondence between string states and Schwarzschild black holes was studied by Horowitz and Polchinski in ref. [39]. Further discussion on the entropy problem from the viewpoint of Matrix theory and the relation between the space-time uncertainty relation and black-hole physics is the subject of later sections.

### 2.3 The space-time uncertainty relation for D-particles

One of the puzzles in proposing the space-time uncertainty relation is how to probe the region  $\Delta X \sim 0$  using strings. Since this leads to large times  $\Delta T \rightarrow \infty$ , the original interpretation was that it provides an explanation why the asymptotic states of any string states, in spite of their intrinsic spatial extension, can be treated as the *local* space-time fields, as in the nonlinear  $\sigma$  model approach to the computation of string S-matrices. Because of this property, the vertex operator of graviton is exactly given by the space-time energy-momentum tensor of strings and hence ensures the equivalence principle. It is, however, desirable to probe the region of short spatial scales directly.

The D-branes, in particular, the point-like D-brane (D-particle), turn out to be ideal probes for that purpose. A D-particle in type IIA superstring theory is characterized by its Ramond-Ramond charge associated with the 1-form RR gauge field, which is identified

---

<sup>||</sup>This argument was given, independently of ref. [39], in a lecture at the 96 Japanese GRG workshop by one of the present author [37].

with the Kaluza-Klein momentum in the 11th direction with compactified radius  $R \sim g_s \ell_s$ . From 10 dimensional viewpoint, it simply behaves as a point particle with mass  $1/g_s \ell_s$ , interacting with each other and with closed strings through open strings whose ends are attached to D-particles. If we apply the space-time uncertainty relation to the open strings between D-particles, we expect that the uncertainty relation holds between the time scale  $\Delta T$  and the spatial scale  $\Delta X$  which is transverse to D-particles. As shown in [27], we can readily derive the characteristic spatial and temporal scales using this relation. Let us repeat the simple argument here.

Consider the scattering of two D-particles of mass  $1/g_s \ell_s$  with the impact parameter of order  $\Delta X$  and the relative velocity  $v$  which is assumed to be much smaller than the light velocity. Then the characteristic interaction time  $\Delta T$  is of order  $\frac{\Delta X}{v}$ . Since the impact parameter is of the same order as the longitudinal length of the open strings mediating the interaction of the D-particles, we can use the space-time uncertainty relation in the form

$$\Delta T \Delta X \sim \ell_s^2 \Rightarrow \frac{(\Delta X)^2}{v} \sim \ell_s^2$$

This gives the order of the magnitude for the minimum possible distances probed by the D-particle scatterings with velocity  $v (\ll 1)$ .

$$\Delta X \sim \sqrt{v} \ell_s. \quad (2.8)$$

To probe short spatial distances, we have to use D-particles with small velocity. However, the slower the velocity is and hence the longer the interaction time is, the larger is the spreading of the wave packet.

$$\Delta X_w \sim \Delta T \Delta_w v \sim \frac{g_s}{v} \ell_s \quad (2.9)$$

since the ordinary time-energy uncertainty relation says that the uncertainty of the velocity is of order  $\Delta_w v \sim g_s v^{-1/2}$  for the time interval of order  $\Delta T \sim v^{-1/2} \ell_s$ . Combining these two conditions, we see that the shortest spatial length is given by

$$\Delta X \sim g_s^{1/3} \ell_s \quad (2.10)$$

and the associated time scale is

$$\Delta T \sim g_s^{-1/3} \ell_s. \quad (2.11)$$

(2.10) is of course the 11 dimensional Planck length which is the characteristic length of M-theory which was first derived in the super YM context in [38]. As argued in ref. [27], it is actually possible to probe shorter lengths than the Planck length if we consider a D-particle in the presence of many ( $=N$ ) coincident D4-branes.

Above derivation of the Planck length emphasizes its meaning as the minimal *observable* distance in a dynamic process of D-particles. We can also derive the same length scale from a slightly different viewpoint of the distance scale below which we cannot discriminate the quantum state of D-particles. For a given velocity  $v$  of a D-particle, the minimal possible distance among D-particles is

$$\Delta X \geq \sqrt{v} \ell_s. \quad (2.12)$$

The ordinary Heisenberg relation  $\Delta p \Delta X \geq 1$ , on the other hand, gives a lower bound for the velocity

$$v \geq \frac{g_s \ell_s}{\Delta X} \quad (2.13)$$

which, by combining with (2.12), leads again to the minimum distance (2.10). Thus, the space-time uncertainty relation clearly says that for each state of a D-particle no information can be stored below the Planck distance and the minimum bit of information carried by a D-particle is given by the Planck volume  $V_P \sim \ell_P^9$  in the transverse space. This is certainly consistent with the holography principle [40] in M theory interpretation of D-particles; however, it does *not* yet prove that the latter is a direct consequence of the space-time uncertainty principle, since we have to actually deal with many-body dynamics of D-particles. Further discussion on the possible relation of the space-time uncertainty relation and the holography will be given in section 4 after summarizing, in section 3, the recent developments in black hole physics from the viewpoint of Matrix theory.

#### 2.4 *The scale symmetry of the effective Super Yang-Mills theory and the space-time uncertainty relation*

Let us now see how the space-time uncertainty principle is encoded in the effective (super) Yang-Mills theory of D-branes. For definiteness, we first concentrate to the case of D-particle and then indicate the generalization to general D-branes only briefly.

The collective variables of  $N$  D-particles are the one-dimensional  $N \times N$  Hermitian matrix field  $X_i(t)$  and their fermionic partners, where  $t$  is the time along the world line of the D-particle.

$$S = \int dt \text{Tr} \left( \frac{1}{2g_s \ell_s} D_t X_i D_t X_i + i\theta D_t \theta + \frac{1}{4g_s \ell_s^5} [X_i, X_j]^2 - \frac{1}{\ell_s^2} \theta \Gamma_i [\theta, X_i] \right) \quad (2.14)$$

where the covariant derivative is defined by  $D_t X = \frac{\partial}{\partial t} X + [A, X]$  with  $A$  is the  $U(N)$  gauge field. This action is invariant under the scale transformation

$$X_i(t) \rightarrow X'_i(t') = \lambda X_i(t), \quad A(t) \rightarrow A'(t') = \lambda A(t), \quad t \rightarrow t' = \lambda^{-1} t \quad (2.15)$$

$$g_s \rightarrow g'_s = \lambda^3 g_s. \quad (2.16)$$

Note the opposite scalings for transverse coordinate  $X_i$  and the time in (2.15). The scaling of the coupling constant (2.16) leads to the characteristic spatial and temporal scales given before on the basis of the space-time uncertainty relation. Thus the effective super Yang-Mills theory naturally encodes the space-time uncertainty principle. This scaling property can be readily extended to general Dp-brane:

$$g_s \rightarrow g'_s = \lambda^{3-p} g_s \quad (2.17)$$

with the same transformation for the transverse coordinate and time as (2.15). The uncertainty relation for D-branes in general says that the long distance phenomena in the (transverse) target space is dual to the short distance phenomena in the world volume and *vice versa*. In particular, for D3-brane ( $p = 3$ ) where the string coupling is invariant, the Yang-Mills theory is invariant corresponding to the scale invariance of the 3+1 dimensional Yang-Mills theory. This means that the dynamics of D3-brane involves all the scales in both the target and world volume, keeping the dual nature of them. For other “dilaton” branes, on the other hand, the effective scales are fixed by the vacuum expectation values of the dilaton. We emphasize that the space-time uncertainty relation itself is valid for general D-branes, irrespectively of such specialities.

In the new interpretation of super Yang-Mills theory as the effective collective field theory of D-branes, the participation of the string coupling in the scale symmetry is allowed, if we remember that the string coupling constant  $g_s$  can in principle be treated

as a dynamical variable corresponding to the vacuum expectation value of the dilaton. Although in the present formulation of the strings and D-branes, the string coupling appears as an external parameter, we should expect that the string coupling should be eliminated from the fundamental (hopefully background independent) formulation of the theory. For example, in the string field formalism, we can indeed eliminate the string coupling by making a shift for the string field as shown in [41].

Now the scaling property constrains the effective action for D-particles with respect to a double power series expansion in  $1/r$  (inverse distances) and  $v$  (velocities), if we neglect the second and higher time derivatives, as

$$S^{eff} = \int dt \sum_{L=0}^{\infty} \sum_{A+2B=L, B \geq 0} R^{3L-3+A+B} \ell_P^{-6L+6-3(A+B)} \times r^A v^B c_{A,B} \quad (2.18)$$

where  $R = g_s \ell_s$ ,  $\ell_P = g_s^{1/3} \ell_s$  are the compactification radius and the Planck length of M-theory, respectively, and  $L$  denotes the number of loops. Here,  $r$  and  $v$  collectively represent the distances and velocities, respectively, of many D-particle systems. The numerical coefficients  $c_{A,B}$  depends on the number  $N$  of D-particles as

$$c_{A,B} = N^{L+1} \sum_{0 \leq n \leq L/2} c_{A,B,n} N^{-2n}.$$

In terms of the string length and the string coupling, the expansion is a double expansion

$$\frac{1}{g_s \ell_s} \left( \frac{g_s \ell_s}{r^3} \right)^{L-4/3} \left( \frac{\ell_s^2 v}{r^2} \right)^B. \quad (2.19)$$

Note that the characteristic combination  $\ell_s^2 v / r^2$  just reflects the space-time uncertainty relation (2.8) for the velocity and the distances, while the combination  $g_s \ell_s / r^3$  reflects the characteristic spatial scale (2.10). Thus the space-time uncertainty relation naturally explains the emergence of the two characteristic quantities by which we can measure the effect of string extension and string coupling in the effective super Yang-Mills model. Of course, the power series expansion loses its meaning when these quantities becomes of order one. We mention that the emergence of the characteristic spatial scale (2.10) is also observed [43] in a different version of Matrix theory, the so-called Matrix string theory [44].

---

<sup>\*\*</sup> This form is first pointed out in ref. [42].



We also note that in terms of the 11 dimensional Newton constant  $G_{11} = g_s^3 \ell_s^9$ , these quantities become ,with  $B = 2n$ ,

$$(\frac{\ell_s^2 v}{r^2})^B \rightarrow \frac{G_{11}^{2n/3}}{R^{2n}} (\frac{v^2}{r^4})^n, \quad (\frac{g_s^{1/3} \ell_s}{r})^{3L} \rightarrow G_{11}^{L/3} r^{-3L}.$$

The classical supergravity contribution corresponds to the "diagonal" terms  $L + 1 = n$ ,  $v^{2(L+1)}/r^{7L}$ . It seems reasonable to interpret the other terms with  $L > n - 1$  or  $n > L + 1$  as the quantum gravity and stringy corrections, respectively. The diagonal contributions have been shown to precisely agree with classical 11 dimensional supergravity up to two loop approximation for 2 [45][42] and 3 body [46] interactions <sup>††</sup> of D-particles for finite  $N$ . Supersymmetry shows that there is no correction for  $n = 0, 1$  terms to the classical contributions. Explicit computations in these references further showed that the correction to the  $n = 2$  term also vanish up to the two-loop approximations, confirming the supersymmetric non-renormalization theorem (see, e. g., [48]).

### 2.5 The Maldacena conjecture and the space-time uncertainty principle

If the above scaling property is interpreted as the symmetry of the theory, it is natural to ask whether the scaling symmetry can be generalized to full conformal symmetry, since we know that the scaling symmetry of the D3-brane is embedded in the full (super) conformal symmetry of the theory. Because of the existence of the maximal supersymmetry, we expect that the full conformal symmetry is unbroken and hence puts stronger constraints on the dynamics of D-particles than the simple scaling arguments. In a recent work [49], it was shown that there exists such a generalization. The D-particle action is indeed invariant under the infinitesimal coordinate transformation

$$\delta_K X_i = 2t X_i, \quad \delta_K A = 2t A, \quad \delta_K t = -t^2, \quad \delta_K g_s = 6t g_s. \quad (2.20)$$

With the scaling symmetry  $\delta_D$  and the trivial time translation symmetry  $\delta_H$ , we have the 1+1 dimensional conformal algebra

$$[\delta_D, \delta_H] = \delta_H, \quad [\delta_D, \delta_K] = -\delta_K, \quad [\delta_H, \delta_K] = 2\delta_D. \quad (2.21)$$

---

<sup>††</sup> There has been some controversy in the case of 3-body interactions, since the first paper in ref. [47] on this. Ref. [46] however showed that the precise agreement holds if we take into account all the relevant terms both in 11D supergravity and Matrix theory, except the recoil effect. Since in this classical approximation, the effective actions in both sides are identical, the recoil effect should also agree.

This is important in view of the conjecture made by Maldacena [50], who proposed essentially that the correspondence between the super Yang-Mills theory and the supergravity implied by the nature of the former as the effective description of the string theory becomes exact in the limit where the higher-excitation modes of open strings are decoupled in the former and the strong curvature and quantum gravity effects are neglected in the latter. The first condition leads us to consider the near horizon region of the D-branes, while the second condition requires the large  $N$  limit with fixed but large  $g_s N \sim g_{\text{YM}}^2 N$ . In the particular case of D3-brane, the near horizon geometry is described by the Anti de Sitter geometry in 4+1 dimensions,  $AdS_5$ . It was then proposed [51][52] to interpret the 3+1 dimensional super Yang-Mills theory as the boundary conformal field theory corresponding to type IIB supergravity in the background of the conformal symmetric  $AdS_5$ .

In ref. [49], it was shown that there exists a similar conformal symmetry for the background fields of D-particle. For example, the 10 dimensional (string frame) metric of  $N$  coincident D-particles are given by

$$ds_{10}^2 = -e^{-2\tilde{\phi}/3} dt^2 + e^{2\tilde{\phi}/3} dx_i^2. \quad (2.22)$$

where

$$e^\phi = g_s e^{\tilde{\phi}}, \quad e^{\tilde{\phi}} = \left(1 + \frac{q}{r^7}\right)^{3/4} \quad (2.23)$$

with  $q = 60\pi^3(\alpha')^{7/2}g_s N$ . If we take the decoupling limit  $\alpha' \rightarrow 0$  keeping

$$U \equiv \frac{r}{\alpha'}, \quad g_{YM}^2 \equiv \frac{g_s}{\alpha'^{3/2}} \quad (2.24)$$

fixed, the metric is reduced to the form

$$ds_{10}^2 = \alpha' \left( -\frac{U^{7/2}}{\sqrt{Q}} dt^2 + \frac{\sqrt{Q}}{U^{7/2}} (dU^2 + U^2 d\Omega_8^2) \right). \quad (2.25)$$

It is easy to check that the metric and the dilaton is invariant under the same scale transformation as before

$$U \rightarrow \lambda U, \quad t \rightarrow \lambda^{-1} t, \quad g_s \rightarrow \lambda^3 g_s, \quad (2.26)$$

Furthermore, they are also invariant under the infinitesimal transformation

$$\delta_K t = -(t^2 + k \frac{g_{YM}^2}{U^5}), \quad (2.27)$$

$$\delta_K U = 2tU, \quad (2.28)$$

$$\delta_K g_s = 6tg_s, \quad (2.29)$$

where  $k$  is a constant independent of the string coupling  $k = 24\pi^3 N$ . Together with the scale and time translation symmetry, they form the same conformal algebra as for the super Yang-Mills quantum mechanics. The appearance of the *field dependent* term, second term in (2.28), is similar to the case of D3 brane. It is now very natural to suppose the validity of the Maldacena conjecture for D-particle case, as in the same sense as for the D3-brane. Namely, the effective super YM quantum mechanics may be interpreted as the *boundary* quantum theory of the D0 metric. For more details on this we refer the reader to [49]. Here, however, we want to emphasize that there is an unsolved question in Maldacena conjecture in general. The question is how to derive the field dependent transformation laws in the bulk from the linear transformation laws in the boundary field theory.

It should be noted that the scaling symmetry naturally leads to the structure which is consistent with the space-time uncertainty relation  $\Delta U \Delta t \sim 1$  for the coordinates  $U$  and  $t$  just as  $\Delta X \Delta T \sim \alpha'$  for the small proper distance  $\Delta X \sim \sqrt{\alpha' Q^{1/2} U^{-7/2}} \Delta U$  and proper time  $\Delta T \sim \sqrt{\alpha' U^{7/2} Q^{-1/2}} \Delta t$ , the latter of which vanishes, for fixed coordinate uncertainties, in the near core limit  $U \rightarrow 0$  while for large  $U$  the situation becomes opposite  $\Delta X \rightarrow 0$ . We also want to emphasize that the Maldacena's conjecture can be regarded as a special case of the original  $s$ - $t$  duality between closed and open strings. Therefore, its consistency with the space-time uncertainty relation and the appearance of the conformal symmetry here again are not at all accident.

It seems natural to interpret the conformal symmetry structure as a mathematical realization of the space-time uncertainty principle, being the transformation groups which leave the form of the super Yang-Mills action invariant. The maximal supersymmetry is crucial here in order to preserve the classical scale and conformal structure in quantum theory. The conformal symmetry in the form (2.27) is then very powerful, since it uniquely determines, combined with the supersymmetric non-renormalization theorem, the classical contribution, i.e. the  $L = n + 1$  terms, of the effective action to all orders in the velocity expansion as shown in [49]. The resulting effective action precisely coincides with the

effective action obtained [42] from the so-called discrete light-cone interpretation of the 11 dimensional supergravity. Hence it supports the interpretation of the effective super Yang-Mills theory of D-particles as the non-perturbative formulation of M theory in the discrete light-cone quantization scheme. Not only that, our arguments almost indicate that the most essential dynamics of D-particles are encapsulated in the space-time uncertainty principle as expressed in the form of the generalized conformal symmetry. Unfortunately, however, our discussion is yet confined to a very special background configurations of D-particles. Therefore, one of the most important direction from this viewpoint would be to generalize the arguments to arbitrary backgrounds.

These structure is not special for D-particle and D3-brane. At least formally, we can extend the conformal symmetry and its consequences to arbitrary  $p$  as will be discussed in [53].

## 2.6 Type IIB matrix model and the space-time uncertainty principle

The D-particle is the lowest dimensional objects in type IIA string theory and M-theory. On the other hand, in type IIB theory, the objects of lowest dimensionality are D-instantons, namely  $p = -1$  D-brane. For the D-instanton, all of the space-time directions, including even the time direction, becomes the transverse directions. The interaction of the D-instantons are therefore described by the open strings which obey the Dirichlet boundary condition with respect to all the space-time directions. Hence the effective action for D-instantons, at least for low-energy phenomena, are described by the 10 dimensional super Yang-Mills theory reduced to a space-time point:

$$S_{\text{eff}} = -\frac{1}{g_s} \text{Tr}([X_\mu, X_\nu])^2 + \text{fermionic part} \quad (2.30)$$

which is covariant from the beginning. We here suppress the string scale by taking the unit  $\ell_s = 1$ .

The dynamics of D-instantons is very special in the sense that the only “physical” degrees of freedom are the lowest massless modes of the open strings. The reason is that only on-shell states satisfying the Virasoro condition are the zero-momentum “discrete” states of the Yang-Mills gauge fields which are identified with  $X_\mu$ ’s and their super partners. The infinite number of the massive degrees of freedom of the open strings should

be regarded as a sort of auxiliary fields. We then expect that the exact effective action for them should be expressible solely in terms of the collective fields which are nothing but the lowest modes  $X_\mu$ . If we further assume that all the higher dimensional objects including the fundamental strings could be various kinds of bound states or collective modes composed of infinitely many D-instantons, we may proceed to postulate that a matrix model defined at a single point might give a possible nonperturbative definition of the type IIB string theory.

This is a very bold assumption: For example, unlike the case of the M(atric) theory, here there is no rationale why we can neglect the configuration in which both instantons and anti-instantons are present. We can only hope that suitable choice of the model might take into account such effect implicitly. A concrete proposal along this line was put forward first by Ishibashi, Kawai, Kitazawa and Tsuchiya [18]. Their proposal is essentially to regard the above simple action as the exact action in the following form.

$$Z_{\text{IKKT}} = \sum_N \left( \prod_\mu \int d^{N^2} X_\mu \right) d^{16} \psi \exp S[X, \psi, \alpha, \beta] \quad (2.31)$$

$$S[X, \psi, \alpha, \beta] \equiv -\alpha N + \beta \text{Tr}_N \frac{1}{2} [X_\mu, Y_\nu]^2 - \dots \quad (2.32)$$

with  $\alpha \propto \frac{1}{g_s \ell_s^4}$ ,  $\beta \propto \frac{1}{g_s}$  and the summation over  $N$  is assumed. Indeed, it has been shown that the model is consistent with the leading long-distance behavior of D-brane interactions expected from the low-energy effective theory, the type IIB supergravity in 10 dimensions.

Is there any signature of the space-time uncertainty principle in this model? Clearly, we cannot extend the previous scaling arguments to the present model, since there is no world volume coordinate. The crucial difference is that now the time is also treated as a matrix. Formally, the type IIB matrix action can be regarded as the T-dual correspondent to the type IIA matrix action. Hence, the scaling of the time matrix is now reversed from the time coordinate of the latter. Thus the scaling property simply leads to the 10 dimensional Planck length  $g_s^{1/4} \ell_s$  to be the characteristic scale for D-instantons. However, D-instantons themselves can never be used as probes for the space-time structure and thus the physical scales can only be determined by the complicated dynamics of the collective modes of infinitely many systems of D-instantons.

If we remember the analogy with ordinary quantum mechanics, it is now tempting to suppose that, instead of the scaling property, some operator algebra is the right tool of discussing the space-time uncertainty principle in the type IIB model, since all the space-time coordinates are treated as (infinite) dimensional matrices. Such an interpretation is proposed in [54], suggesting that the model should be formulated as a microcanonical statistical system with a constraint on the commutation relations of the coordinate matrices in the form

$$\langle \frac{1}{2}([X_\mu, X_\nu])^2 \rangle \equiv \lim_{N \rightarrow \infty} \frac{1}{N} \text{Tr} \frac{1}{2}([X_\mu, X_\nu])^2 = \alpha'^2. \quad (2.33)$$

which leads to the inequality

$$\sqrt{-\langle ([X^0, X^i])^2 \rangle} \geq \alpha', \quad (2.34)$$

in conformity with the space-time uncertainty relation  $\Delta T \Delta X \geq \alpha'$ . The integration measure is determined to be that of the maximal supersymmetric model, namely the fermionic determinant, if we require the cluster decomposition property of space-time. Because of the equivalence of the canonical and microcanonical ensemble, we can then derive the IKKT model using the standard arguments of statistical mechanics. For more motivations supporting this interpretation and the relation with conformal invariance, we refer the reader to ref. [54].

Now that we have summarized the space-time uncertainty principle in string theory, we again turn to the problem of black holes.

### 3. Matrix black holes

Much progress has been made in understanding microscopic origin of Bekenstein-Hawking entropy in the past few years [14] (for reviews see [15]) from the standpoint of string theory. Since most work were based on the physics of D-branes, it has been hard to go beyond the case of extremal black holes. Matrix theory provides a unique opportunity to understand the standard Schwarzschild black holes, as it purports to provide a nonperturbative formulation for M theory. As we have seen in the last section, it also forms a natural ground for space-time uncertainty relation. We will see in the next section how the two seemingly different issues are naturally tied up together.

By boosting a hole with an extremely large longitudinal momentum, one effectively puts the hole to a background of a large number of D0-branes which are BPS states. Now the properties of the quantum black hole is determined by the statistical mechanics of a large number of D-partons. The geometry is probed by sending in a classical probe which also consists of a large number of D-partons.

The first observation, due to Banks et al. [55], is that for a finite longitudinal cut-off  $R$  and a black hole of radius  $r_s > R$ , it is necessary to boost the hole in order to fit it into the asymptotic box size  $R$ . Asymptotically, one can apply the Lorentz contraction formula  $r_s e^{-\alpha}$ , where  $\alpha$  is the rapidity parameter, roughly equal to  $M/P_-$ . The minimal boost is determined by  $r_s e^{-\alpha} = R$ , or  $P_- = r_s M/R$ . In matrix theory,  $P_- = N/R$ , where  $N$  is the number of partons. The above formula says that  $N = r_s M \sim S$ , where  $S$  is the entropy of the hole. This condition then says that the minimal number of partons required to account for entropy  $S$  is just  $S$ , a physically appealing claim.

Geometrically, one might wonder how the Lorentz contraction could happen to a horizon, since by definition horizon is a null surface which is independent of the coordinates used. Indeed it can be shown that in the boosted frame, the size of horizon remains the same. What the boost does to the black hole is to change the relation between the size of the horizon and the asymptotic radius of the longitudinal direction, if the hole is put on a periodic circle. It can be shown that for the horizon size to be  $r_s$  while the asymptotic box size to be  $R$ , the hole must carry a minimal momentum as determined naively in the last paragraph [56].

Another point we want to emphasize here is that when the size of the hole fits the box size, it looks more like a black string. Indeed, a black string becomes unstable at the special point  $N \sim S$ . Since the horizon area of the black hole of the same size and same momentum is greater than that of the black string when one slightly increases the momentum, the black string will collapse to a black hole.

We will be able to explain the size of the hole and its entropy only up to a numerical coefficient, thus whenever we write down a formula that is valid only up to a numerical coefficient. In D dimensional space-time, the size of the Schwarzschild black hole and its

entropy, written in terms of the mass are given by

$$r_s^{D-3} = G_D M, \quad S = \frac{r_s^{D-2}}{G_D} = G_D^{\frac{1}{D-3}} M^{\frac{D-2}{D-3}}. \quad (3.35)$$

At the special kinetic point  $N \sim S$ , we use the second relation in (3.35) to solve  $M$  in terms of  $N$ :

$$M = G_D^{-\frac{1}{D-2}} N^{\frac{D-3}{D-2}},$$

thus the light-cone energy

$$E_{LC} = R G_D^{-\frac{2}{D-2}} N^{\frac{D-4}{D-2}}, \quad (3.36)$$

and the size of the hole

$$r_s = (G_D N)^{\frac{1}{D-2}}. \quad (3.37)$$

As we argued before, the boosted black hole at the transition point  $N \sim S$  can be either regarded as a black string, if the longitudinal momentum is slightly smaller than the critical value, or a black hole if the longitudinal momentum is slightly larger. In the former case, one needs to excite longitudinal objects such as longitudinally wrapped membrane in matrix theory, thus the momentum modes in the low energy nonabelian field theory are relevant. Actually the hole phase is easier to account for. Only the zero modes, in other words, the motion of D0-branes in the open space are relevant.

When the Born-Oppenheimer approximation is valid, the one-loop, spin-independent potential between two D0-branes is given in [38]. The assumption that the Born-Oppenheimer approximation is valid for a black hole implies that the dominant part of the black hole is a gas of D0-branes, such that for dynamic purposes one can integrate out off-diagonal variables. In  $D$  dimensional space-time, when M theory is compactified on a torus  $T^{11-D}$ , the analogous potential between two D0-branes can be obtained from that in 11D by summing over infinitely many images on the covering space of the torus:

$$L = \frac{1}{2R}(v_1^2 + v_2^2) + \frac{c_D G_D}{R^3} \frac{(v_1 - v_2)^4}{r^{D-4}}, \quad (3.38)$$

if the exchange of supergraviton producing the potential does not cause longitudinal momentum transfer. The above form is certainly appropriate for D0-branes, and for threshold bound states of D0-branes. It is a simple matter to use the above interaction and the Heisenberg uncertainty relation to derive formulas in (3.36) and (3.37) [56][57].



However, it is rather bizarre to consider a black hole whose size is comparable to the size of one spatial dimension of the universe. To lift the restriction, we need to consider a highly boosted black hole such that its longitudinal size is much smaller than its transverse size and has nothing to do with  $R$ . In such a situation, we postulate that the black hole is better viewed as a gas of  $S$  clusters whose average longitudinal momentum is  $p_- = N/(SR)$  [58]. For the time being  $S$  has nothing to do with entropy. The interaction energy for zero longitudinal momentum exchange is given by the above formula with  $1/R^2$  replaced by  $p_-^2$ . Here we need to take one step further, to assume that for processes in which longitudinal momentum transfer occurs the interaction takes the more or less the same form, then the total interaction energy of the gas is

$$V \sim G_D \sum_{\delta p_- = 0}^{N/(SR)} \sum_{a,b} \frac{p_- v^2 p_- v^2}{R r_s^{D-4}}. \quad (3.39)$$

where the maximal longitudinal momentum exchange is limited by  $p_- = N/(SR)$  too.

The velocity of a cluster is bounded by the Heisenberg uncertainty relation  $p_- v \sim 1/r_s$ , thus the interaction energy is bounded by

$$V \sim G_D \frac{N}{S} S^2 \frac{S^2 R}{N^2 r_s^D} \sim E_T \frac{G_D S}{r_s^{D-2}}. \quad (3.40)$$

The virial theorem  $V \sim E_T$  implies that  $S \sim r_s^{D-2}/G_D$ , the Bekenstein formula. Here we assumed that the number of clusters determines entropy.

Note that the above is the *lower* bound for the interaction energy, according to Heisenberg relation, thus the Bekenstein-Hawking formula is a *upper* bound. This explains two interesting points.

1. If the localization size of a cluster is  $r < r_s$ , then we would obtain a formula  $S \sim r_s^{D-2} r^2 / G_D$  which is smaller than the hole entropy.
2. Even when the localization size is  $r_s$ , the Heisenberg relation must be saturated in order to attain the maximal entropy given by the Bekenstein-Hawking formula.

As argued in [59], it is also possible to account for the correct virial theorem using spin-dependent interactions among individual D0-branes.

It can be shown that the contribution to the potential energy from other forms of interaction is the same order as (3.40). The fact that there are about  $S$  clusters suggests that these clusters obey Boltzmann statistics. This is easy to justify for large  $N$ , since each cluster may have some fluctuation in its longitudinal momentum. It is also possible that some background whose kinetic energy is negligible is responsible for the distinguishable clusters.

The correspondence principle of Horowitz and Polchinski, as explained already in the previous section, can be derived, using the above information from matrix theory. Furthermore, the dynamic details of string/hole transition can be studied in the matrix string picture [60]. For related works, see [61].

We now recall what we have learned. Under a large boost, the longitudinal size of the black hole as seen by a distant observer contracts enormously. Denote this size by  $r_L$ , then its relation to the Schwarzschild radius is  $r_L = r_s e^{-\alpha} = r_s M/P_-$ . Again to a distant observer, the black hole is described as a gas of clusters each carrying longitudinal momentum  $p_- \sim 1/r_L \sim P_-/(r_s M) = P_-/S$ . This gas is best viewed as a version of the quantum stretched horizon, and the fact that there are  $S$  such clusters strongly suggests a holographic picture, namely the black hole physics is effectively described by clusters in a  $D - 1$  dimensional theory. We shall see in the next section, that this picture also motivates a generalization of space-time uncertainty relation to M theory, whose validity goes beyond string theory.

## 4. Space-time uncertainty relation and holography

### 4.1 *M-theory interpretation of the space-time uncertainty relation*

The space-time uncertainty relation  $\Delta X \Delta T \geq \alpha'$  has a firm ground in string theory, as we have argued in section 2 where its possible relevance as the underlying principle for holography is also suggested. In the context of matrix theory, the space-time uncertainty relation takes the form

$$\Delta X_T \Delta T \geq \frac{l_p^3}{R} \quad (4.41)$$

where  $\Delta X_T$  and  $\Delta T$  are the characteristic scales with respect to spatial transverse distances and time, respectively. This relation is about processes involving individual D-

partons, whose longitudinal wave length is given by  $R$ . Admittedly, this is a rather weak relation in a process when more complicated composite objects are involved.

A reinterpretation of the above relation from the viewpoint of M theory allows us to write down a more general relation. Note that a process involving individual D-partons necessarily smears over the longitudinal direction, thus the uncertainty in this direction  $\Delta X_L = R$  is maximal. Relation (4.41) is rewritten as

$$\Delta X_T \Delta X_L \Delta T \geq l_p^3, \quad (4.42)$$

this relation <sup>‡‡</sup> refers only to the fundamental length scale in M theory, the Planck length, thus it is a natural candidate for the generalized uncertainty relation in M theory. In the previous section, we learned that in a highly boosted object the natural constituent is a cluster whose longitudinal wave length does not exceed the longitudinal size of the macroscopic object. We now argue that relation (4.42) is the correct relation for a process involving such a cluster. It is trivially true for threshold bound state, since it is just a boosted parton and according to Lorentz invariance  $\Delta X_L$  contracts, while  $\Delta T$  is dilated by a same factor. An object carrying the same amount of longitudinal momentum can be regarded as an excited state of the threshold bound state, therefore intuitively as a probe it can not probe a transverse distance shorter than a threshold bound state can do. Thus, relation (4.42) must also hold for such a probe. Note also that this relation is Lorentz invariant.

A good example of a cluster is the one discussed in [56]. That is a spinning membrane. If we set its transverse size  $X_T = 1$ , then the time scale is  $T = N/R$ . Now since the spinning membrane carries longitudinal momentum  $P_- = N/R$  and so its longitudinal extension is  $X_L = 1/p_- = R/N$ . The cubic uncertainty relation is satisfied.

As explained in the last part of section 2, one of us argued in [54] that the string uncertainty relation serves as a guiding principle in constructing the IIB matrix model. We suspect that if the M theory uncertainty relation (4.42) has any deep meaning, it may also underlie the ultimate nonperturbative and covariant formulation of M theory whose validity goes beyond Matrix theory which is only formulated in the light- cone frame (or IMF).

---

<sup>‡‡</sup> This is reminiscent of a triple uncertainty relation  $\Delta X \Delta Y^2 \geq g_s \ell_s^3$  suggested in [27].

In the AdS/CFT correspondence conjectured by Maldacena [50], the holographic bound is satisfied provided one notices the UV/IR correspondence between a bulk process and a process in the boundary theory. To see how our uncertainty relation (4.42) fits into this context, we examine the  $AdS_7 \times S^4$  case. The reason for picking out this one is obvious: There is no underlying string theory, thus relation (4.42) is most suitable. The metric of the  $AdS_7$  is given by

$$ds^2 = \frac{r}{R}(-dt^2 + \sum_{i=1}^5 dx_i^2) + \frac{R^2}{r^2}dr^2, \quad (4.43)$$

where the coordinates  $(t, x_i)$  are the natural ones on fivebranes, we have set  $l_p = 1$ .

The simplest way to probe this background is through the use of open membranes. An open membrane appears as a string inside fivebranes. Our uncertainty relation says that if the uncertainty in the brane theory is given by  $\Delta T$  and  $\Delta X_L$ , then necessarily the membrane state will have an extension in the transverse direction (the radial direction) at least  $\Delta X_T \sim 1/(\Delta T \Delta X_L)$ . This translates into a statement in terms of coordinates in (4.43) that  $\Delta r \sim 1/(\Delta t \Delta x)$ , thanks to the peculiar form of the metric. It is easy to see that this is a generalization of the UV/IR correspondence observed by Susskind and Witten, as discussed in the next subsection.

A further support to the above relation is the scaling argument similar to the one for the original uncertainty relation for strings and D-branes.  $r$  naturally corresponds to scalars on fivebranes. There are 5 scalars on each fivebrane representing transverse fluctuations, in a tensor super-multiplet. Now the scalar in the 6D conformally invariant theory has dimension 2, namely if  $x_i \rightarrow \lambda x_i$ , then  $\phi \rightarrow \lambda^{-2}\phi$ . This is precisely our stated uncertainty relation.

The generalized space-time uncertainty relation says nothing about the nature of the theory living on the fivebranes, except for the above scaling property. As observed in [62], so long if one has a consistent short distance theory on the branes, the theory is able to account the entropy in the whole bulk space. This is certainly a good qualitative argument for Maldacena's conjecture. To obtain the right Bekenstein-Hawking bound in this particular example, we need to use another set of coordinates, similar to those in [62]. As pointed out by Banks, the number of degrees must be proportional to  $N^3$  in this case, in other words, the energy density must be proportional to  $N^3/\delta^6$ ,  $\delta$  is the UV cut-

off. The close relation between holography and spacetime uncertainty relation certainly suggests that the latter should have a deep meaning. Related views on the interpretation of the space-time uncertainty principle and its relation to holography are independently pointed out by Minic [63].

#### 4.2 *The space-time uncertainty relation and the UV/IR correspondence of Susskind-Witten*

Finally, we want to emphasize that the UV/IR correspondence underlying Susskind-Witten's derivation [62] of the holographic bound for the entropy of the d3-brane is regarded as a consequence of the string space-time uncertainty relation. Following [62], let us assume the infrared cutoff of the bulk theory to be  $A^{1/3} \sim R/\delta$  with respect to the volume  $A$  of the boundary. Here we use the corrected metric ( $R = \ell_s(g_s N)^{1/4}$ ) for the  $AdS_5$  space

$$ds^2 = R^2 \left[ \frac{4dx^i dx^i}{(1-r^2)^2} - dt^2 \frac{(1+r^2)^2}{(1-r^2)^2} \right]. \quad (4.44)$$

This cutoff is equivalent to introducing the uncertainty  $\Delta r \sim \delta$  in the radial ( *i.e.*, transverse) direction orthogonal to the boundary  $r = 1$ . In terms of the proper distance,  $\Delta X \sim R\delta/\delta = R$ . Then by the string space-time uncertainty relation, we have the corresponding cutoff  $\Delta T \sim \ell_s^2/R$  for the proper time, or, in terms of the coordinate time,  $\Delta t \sim \ell_s^2\delta/R^2$ . If we use the dimensionless unit of ref. [62] for the surface theory, we have  $\Delta t \sim \delta$ . This explains the UV cutoff for the boundary theory adopted in [62], where the coordinates of the boundary theory is assumed to be  $(t, x^i)$  with  $x^i x^i \sim 1$ , which leads to the number of degrees of freedom  $N_{dof} = N^2/\delta^2 = A/G_5$ . It is easy to extend the arguments to M2 and M5 branes by using appropriate coordinates similar to (4.44) and determine the  $N$  dependencies required for agreement with the Beckenstein-Hawking formula. For M5 with  $R \sim \ell_P N^{1/3}$  and M2 with  $R \sim \ell_P N^{1/6}$ , we need  $N_{dof} \sim N^3/\delta^5$  and  $N_{dof} \sim N^{3/2}/\delta^2$ , respectively.

It should be kept in mind that, because of the different choice of the coordinates, the uncertainty relation for the coordinate time and coordinate spatial distance now takes the form  $\Delta t \Delta r \sim \ell_s^2 \delta^2 / R^2$  which depends on the cutoff  $\delta$ . In the metric of the type (4.43) (or (2.25) for D0 case), on the other hand, the (generalized) uncertainty relation is valid in the same forms for the proper distances and the coordinate distances. For other choices

of coordinates, we must be careful in applying the uncertainty relation. In view of the original derivation of the space-time uncertainty relation, as explained in subsection 2.1, we have to use the proper distances for physical space-time lengths,  $\Delta X$  and  $\Delta T$ .

## 5. Conclusion

We have reviewed the present status of string theory from the viewpoint of the short-distance space-time structure and black-hole entropy. In particular, we have mainly emphasized the space-time uncertainty relation as a possible guiding principle for understanding some universal short-distance structure in perturbative and non-perturbative string and M theories. We have also suggested the relevance of the space-time uncertainty relation for holography.

There are of course other viewpoints in aiming towards the same goal. Besides some related approaches already mentioned, we have not emphasized, for example, the viewpoint of noncommutative geometry as a possible mathematical framework for formulating the noncommutative structure of space-time geometry at short distances. For an example of such approaches, we refer the reader to [64][65] and to different attempts such as [66][67]. We hope that our physical and intuitive viewpoint would play a useful role which is complementary to more abstract and formal investigations. In particular, we expect that one of the most intriguing questions is how to unify various conformal symmetries, which are now turned out to be the formal bases for the space-time uncertainty principle, and how to incorporate them into a fundamental mathematical formalism.

### *Acknowledgments*

In preparing the present article, we have benefited from conversations and e-mail exchanges with A. Jevicki, M. Kato, Y. Kazama, H. Kawai, E. Martinec, D. Minic. The work of M. L. is supported by DOE grant DE-FG02-90ER-40560 and NSF grant PHY 91-23780. The work of T. Y. is supported in part by Grant-in-Aid for Scientific Research (No. 09640337) and Grant-in-Aid for International Scientific Research (Joint Research, No. 10044061) from the Ministry of Education, Science and Culture.

## References

- [1] G. Veneziano, *Nuovo Cim.* 57A(1968) 190.
- [2] We want to emphasize that the bibliography of the present paper is not, in any sense, aiming a complete listing of all relevant works. We would like to apologize all authors if their works related to our subjects are not cited here.  
  
For more extensive reviews and references on general string theory, we refer the reader to M. Green, J. Schwarz and E. Witten, *Superstring Theory* Vol. 1 and 2. Cambridge Univ. Press (1987),  
  
J. Schwarz (ed.), *Superstrings: the first 15 years of superstring theory* Vol 1 and 2, World Scientific (1985),  
  
and other review articles referred to below.
- [3] Y. Nambu, *Proc. Intl. Conf. on Symmetries and Quark Models* (Wayne state univ.)(1969).  
H. B. Nielsen, *15th Conf. on High Energy Physics* (Kiev)(1970).  
L. Susskind, *Nuovo Cim.* 69 (1970) 457.
- [4] J. A. Shapiro, *Phys. Rev. D* 11(1975) 2937.
- [5] T. Yoneya, *Prog. Theor. Phys.* 51(1974) 1907; *Lett. Nuovo. Cim.* 8(1973)951; *Prog. Theor. Phys.* 56(1976)1310.
- [6] J. Scherk and J. Schwarz, *Nucl. Phys.* B81(1974)118; *Phys. Lett.* 57B(1975)463.
- [7] F. Gliozzi, J. Scherk and D. Olive, *Nucl. Phys.* B122 (1977)253.
- [8] For a review on recent developments, see, e.g.,  
J. Polchinski, *Rev. Mod. Phys.* 68(1996)1245.
- [9] K. Kikkawa and M. Yamasaki, *Phys. Lett.* B149(1984)357.
- [10] A. Sen, *Int. Jour. Mod. Phys.* A9(1994)3707.
- [11] E. Witten, *Nucl. Phys.* B443(1995)85.
- [12] J. Polchinski, *Phys. Rev. Lett.* 75(1995)4724.
- [13] G. Horowitz and A. Strominger, *Nucl. Phys.* B360 (1991)197.
- [14] A. Strominger and C. Vafa, *Phys. Lett.* B379(1996)99.
- [15] For reviews, see, e.g.,  
J. Maldacena, hep-th/9607235.  
A. W. Peet, hep-th/9712253.
- [16] E. Witten, *Nucl. Phys.* B460(1996)335.

- [17] T. Banks, W. Fischler, S. H. Shenker and L. Susskind, Phys. Rev. D55(1997)5112.
- [18] N. Ishibashi, H. Kawai, Y. Kitazawa and A. Tsuchiya, Nucl. Phys B498(1997)467.
- [19] L. Susskind, hep-th/9704080.
- [20] N. Seiberg, Phys. Rev. Lett. 79(1997) 3577.  
See also, A. Sen, hep-th/9709220.
- [21] For a previous review on space-time uncertainty principle, see  
T. Yoneya, hep-th/9707002 (to appear in the Proceedings of the APCPT-ICTP Joint International Conference '97).
- [22] T. Yoneya, *Duality and Indeterminacy Principle in String Theory* in "Wandering in the Fields", eds. K. Kawarabayashi and A. Ukawa (World Scientific, 1987), p. 419: see also *String Theory and Quantum Gravity* in "Quantum String Theory", eds. N. Kawamoto and T. Kugo (Springer, 1988), p. 23.
- [23] D. Amati, M. Ciafaloni and G. Veneziano, Phys. Lett. 197B, 81(1987).
- [24] D. Gross and P. Mende, Nucl. Phys. B303, 407(1988).
- [25] For a review and a list of references related to this subject, see, e. g., L. J. Garay, Int. J. Mod. Phys. A10, 145(1995). For other recent attempts, see also G. Amelio-Camelin, J. Ellis, N. E. Mavromatos and D. V. Nanopoulos, hep-th/9701144 and references therein.
- [26] T. Yoneya, Mod. Phys. Lett. A4, 1587(1989).
- [27] M. Li and T. Yoneya, Phys. Rev. Lett. 78 (1997) 1219.
- [28] J. J. Atick and E. Witten, Nucl. Phys. **B310**, 291(1988).
- [29] See, e.g., S. Doplicher, K. Fredenhagen and J. E. Roberts, Commun. Math. Phys. 172 (1995) 187; Phys. Lett. B331 (1994) 39, and references therein.
- [30] A. Jevicki, M. Li and T. Yoneya, Nucl. Phys. B448(1995) 277.
- [31] I. Klebanov, Proceedings of the Trieste Spring School 1991, eds. J. Harvey et al. (World Scientific, Singapore, 1991);  
A. Jevicki, hep-th/9309115;  
J. Polchinski, hep-th/9411028.
- [32] S. Shenker, hep-th/9509132.
- [33] L. Susskind, Phys. Rev. D49 (1994) 6606.
- [34] A. Jevicki and T. Yoneya, Nucl. Phys. B411(1994) 64.
- [35] T. Yoneya, p. 178 in "Frontiers in Quantum Field Theories" (World Scientific, 1996), hep-th/9603096.



- [36] For review, see, e.g.,  
L. Susskind, hep-th/9511227; D. Bigatti and L. Susskind, hep-th/9712072.
- [37] T. Yoneya, Proc. of the Sixth Workshop on General Relativity and Gravitation (Dec. 2-5, 1996).
- [38] M. Douglas, D. Kabat, P. Pouliot and S. Shenker, Nucl. Phys. B485(1997)85.
- [39] G. Horowitz and J. Polchinski, Phys. Rev. D55(1977)6189.
- [40] G. 'tHooft, *Dimensional Reduction in Quantum Gravity*, gr-qc/9310026; L. Susskind, J. Math. Phys. 36 (1995)6377.
- [41] T. Yoneya, Phys. Lett. 197B(1987) 76.  
R. C. Myers, S. Penati, M. Pernici, and A. Strominger, Nucl. Phys. B310(1988)25.  
H. Hata and Y. Nagoshi, Prog. Theor. Phys. 80(1988) 80.  
O. Bergman and B. Zwiebach, Nucl. Phys. B441 (1995) 441.
- [42] K. Becker, M. Becker, J. Polchinski and A. Tseytlin, Phys. Rev. D56(1997) 3174.
- [43] S. B. Giddings, F. Hacqueboard and H. Verlinde, hep-th/9804121.
- [44] R. Dijkgraaf, E. Verlinde and H. Verlinde, Nucl. Phys. B500(1997)43.
- [45] K. Becker and M. Becker, Nucl. Phys. B506(1997)48.
- [46] Y. Okawa and T. Yoneya, hep-th/9806108.
- [47] M. Dine and A. Rajaraman, hep-th/9710174.  
M. Fabbri, G. Ferretti and R. Iengo, hep-th/9806018, hep-th/9806166.  
W. Taylor IV and M. Van Raamsdonk, hep-th/9806066.  
R. Echols and J. P. Gray, hep-th/9806109.  
J. McCarthy, L. Susskind and A. Wilkins, hep-th/9806136.
- [48] S. Paban, S. Sethi and M. Stern, hep-th/9805018.  
See also, M. Dine, R. Echols and J. P. Gray, hep-th/9805007.
- [49] A. Jevicki and T. Yoneya, hep-th/9805069.
- [50] J. Maldacena, hep-th/9711200.  
N. Itzhaki, J. Maldacena, J. Sonnenschein and S. Yankielowicz, hep-th/9802042.  
See also earlier works,  
S. Gubser, I. Klebanov and A. Peet, Phys. Rev. D54(1996) 3916.  
I. Klebanov, Nucl. Phys. B496 (1997) 231.  
S. Gubser, I. Klebanov and A. Tseytlin, Nucl. Phys. B499 (1997) 217.
- [51] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, hep-th/9802109.

- [52] E. Witten, hep-th/9802150.
- [53] A. Jevicki, Y. Kazama and T. Yoneya, to appear.
- [54] T. Yoneya, Prog. Theor. Phys. 97(1997) 949.
- [55] T. Banks, Fischler, I. R. Klebanov and L. Susskind, Phys. Rev. Lett. 80(1997)226.
- [56] G. Horowitz and E. Martinec, Phys. Rev. D57(1998)4935.
- [57] T. Banks, Fischler, I. R. Klebanov and L. Susskind, hep-th/9711005.
- [58] M. Li and E. Martinec, hep-th/9801070.
- [59] M. Li, hep-th/9710226.
- [60] M. Li, E. Martinec and V. Sahakian, to appear.
- [61] H. Liu and A. A. Tseytlin, hep-th/9712063; D. Lowe, hep-th/9802173; D. Kabat and G. Lifschytz, hep-th/9806214.
- [62] L. Susskind and E. Witten, hep-th/9805114.
- [63] D. Minic, private communication.
- [64] M. Douglas, hep-th/9610041.
- [65] A. Conne, M. Douglas and A. Schwarz, hep-th/9711262.
- [66] S. Hirano and M. Kato, Prog. Theor. Phys. 98(1997) 1371.
- [67] I. Oda, Nucl. Phys. B516(1998)160.